Global Wild Poliovirus 2014 - 2019



Country or territory ³	Wild virus type 1 confirmed cases								Wild virus type 1 reported from other sources ²							
	Full year total					01 Jan - 12 Nov ¹		Date of most	Full year total					01 Jan - 12 Nov ¹		Date of most
	2014	2015	2016	2017	2018	2018	2019	recent case	2014	2015	2016	2017	2018	2018	2019	recent virus
Pakistan	306	54	20	8	12	8	82	21-Oct-19	127	84	62	110	141	102	289	25-Oct-19
Afghanistan	28	20	13	14	21	19	20	28-Sep-19	17	20	2	42	83	51	43	24-Sep-19
Nigeria	6	0	4	0	0	0	0	21-Aug-16	1		1 ⁶					27-Sep-16
Iran	0	0	0	0	0	0	0	NA							3	20-May-19
Israel ⁴	0	0	0	0	0	0	0	NA	14							30-Mar-14
West Bank and Gaza	0	0	0	0	0	0	0	NA	1							05-Jan-14
Somalia	5	0	0	0	0	0	0	11-Aug-14								
Cameroon	5	0	0	0	0	0	0	09-Jul-14								
Equatorial Guinea	5	0	0	0	0	0	0	03-May-14								
Iraq	2	0	0	0	0	0	0	07-Apr-14								
Syrian Arab Republic	1	0	0	0	0	0	0	21-Jan-14								
Ethiopia	1	0	0	0	0	0	0	05-Jan-14								
Total	359	74	37	22	33	27	102		160	104	65	152	224	153	335	
Total wild virus type 1	359	74	37	22	33	27	102									
Total wild virus type 3	0	0	0	0	0	0	0									
Tot. in endemic countries	340	74	37	22	33	27	102									
Tot. in non-end countries	19	0	0	0	0	0	0									
No. of countries (infected)	9	2	3	2	2	1	0									
No. of countries (endemic)	3	25	25	3	3	1	0									

Countries in yellow are endemic. ¹Data reported to WHO HQ on 13 Nov 2018 for 2018 data and 12 Nov 2019 for 2019 data.

⁵Between 27 Sep 2015 and 27 Sep 2016, Nigeria was not classified as endemic. NA - Most recent case had onset prior to 1999. ⁶Exceptionally reporting case-contact of a positive index case given the date of collection is later than the onset date of the most recent WPV.

²Wild viruses from environmental samples, selected contacts, healthy children and other sources. Last WPV type 3 had its onset on 10 November 2012. ³In March 2014, a serotype 1 wild poliovirus was detected in an environment specimen from Brazil, further investigation indicates this is an isolated event without evidence of circulation. ⁴Results are based on L20B positive culture. Prior to reporting week 16, 2014, results were based on a combination of direct qRT-PCR on RNA from concentrated sewage and L20B positive culture.