

Global Wild Poliovirus 2015 - 2020

| Country or territory ³ | Wild virus type 1 confirmed cases | | | | | | | | Wild virus type 1 reported from other sources ² | | | | | | | |
|-----------------------------------|-----------------------------------|----------------|----------------|------|------|-----------------------------|------|--------------|--|----------------|------|------|------|------------------------------|------|--------------|
| | Full year total | | | | | 01 Jan -18 Feb ¹ | | Date of most | Full year total | | | | | 01 Jan - 18 Feb ¹ | | Date of most |
| | 2015 | 2016 | 2017 | 2018 | 2019 | 2019 | 2020 | recent case | 2015 | 2016 | 2017 | 2018 | 2019 | 2019 | 2020 | recent virus |
| Pakistan | 54 | 20 | 8 | 12 | 144 | 4 | 17 | 02-Feb-20 | 84 | 62 | 110 | 141 | 389 | 27 | 29 | 20-Jan-20 |
| Afghanistan | 20 | 13 | 14 | 21 | 29 | 2 | 0 | 07-Dec-19 | 20 | 2 | 42 | 83 | 56 | 9 | 2 | 08-Jan-20 |
| Nigeria ³ | 0 | 4 | 0 | 0 | 0 | 0 | 0 | 21-Aug-16 | | 1 ⁴ | | | | | | 27-Sep-16 |
| Iran | 0 | 0 | 0 | 0 | 0 | 0 | 0 | NA | | | | | 3 | | | 20-May-19 |
| Total (Type1) | 74 | 37 | 22 | 33 | 173 | 6 | 17 | | 104 | 64 | 65 | 224 | 448 | 36 | 31 | |
| Tot. in endemic countries | 74 | 37 | 22 | 33 | 173 | 6 | 17 | | | | | | | | | |
| Tot. in non-end countries | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | | | | | | | | |
| No. of countries (infected) | 2 | 3 | 2 | 2 | 2 | 0 | 1 | | | | | | | | | |
| No. of countries (endemic) | 3 | 2 ³ | 2 ³ | 3 | 3 | 3 | 3 | | | | | | | | | |

Countries in yellow are endemic.¹Data reported to WHO HQ on 19 Feb 2019 for 2019 data and 18 Feb 2020 for 2020 data.

²Wild viruses from environmental samples, selected contacts, healthy children and other sources. Last WPV type 3 had its onset on 10 November 2012. ³Between 27 Sep 2015 and 27 Sep 2016, Nigeria was not classified as endemic. NA - Most recent case had onset prior to 1999. ⁴Exceptionally reporting case-contact of a positive index case given the date of collection is later than the onset date of the most recent WPV.