Global Wild Poliovirus 2016 - 2022



Country or territory ³	Wild virus type 1 confirmed cases										Wild virus type 1 reported from other sources ²							
	Full year total						01 Jan-22 Nov ¹		Date of most recent	Full year total						1 Jan-22 Nov ¹	Date of most recent	
	2016	2017	2018	2019	2020	2021	2021	2022	case	2016	2017	2018	2019	2020	2021	2022	virus	
Afghanistan	13	14	21	29	56	4	1	2	29-Aug-22	2	42	83	60	49	1	11	02-Nov-22	
Pakistan	20	8	12	147	84	1	1	20	15-Sep-22	62	110	141	405	455	65	36	03-Oct-22	
Mozambique	0	0	0	0	0	0	0	8	10-Aug-22									
Malawi	0	0	0	0	0	1	0	0	19-Nov-21									
Iran	0	0	0	0	0	0	0	0	NA				3				20-May-19	
Nigeria ³	4	0	0	0	0	0	0	0	21-Aug-16	14							27-Sep-16	
Total (Type1)	37	22	33	176	140	6	2	30		65	152	224	468	504	66	47		
Tot. in endemic countries	37	22	33	176	140	5	2	22										
Tot. in non-end countries	0	0	0	0	0	1	0	8										
No. of countries (infected)	3	2	2	2	2	3	2	3										
No. of countries (endemic)	2 ³	2 ³	3	3	3	2	2	2										
Total Female	13	7	18	72	59	2	0	9										
Total Male	24	15	15	104	81	4	2	21										

Countries in yellow are endemic. ¹Data reported to WHO HQ on 23 Nov. 2021 for 2021 data and 22 Nov. for 2022 data.

²Wild viruses from environmental samples, selected contacts, healthy children and other sources. Last WPV type 3 had its onset on 10 November 2012. ³Between 27 Sep 2015 and 27 Sep 2016, Nigeria was not classified as endemic, and declared Wild polio-free as of June 2020. ⁴Exceptionally reporting case-contact of a positive index case given the date of collection is later than the onset date of the most recent WPV.